



PATIENT & CAREGIVER EDUCATION

Pazopanib

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Votrient

Brand Names: Canada

PMS-Pazopanib; Votrient

Warning

- Severe and sometimes deadly liver problems have happened with this drug. Call your doctor right away if you have signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.

What is this drug used for?

- It is used to treat kidney cancer.
- It is used to treat soft tissue sarcoma.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor **BEFORE** I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have liver disease or raised liver enzymes.
- If you are being treated with other cancer therapy. The chance of severe and sometimes deadly side effects may be raised.
- If you are taking any of these drugs: Cimetidine, dexamprazole, esomeprazole, famotidine, lansoprazole, nizatidine, omeprazole, pantoprazole, rabeprazole, or ranitidine.
- If you take any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for HIV, infections, seizures, and others. There are many drugs that must not be taken with this drug.
- If you are pregnant or may be pregnant. This drug may cause harm to an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant. If you may become pregnant, you must use birth control while taking this drug and for some time after the last dose. Ask your doctor how long to use birth control. If you get pregnant, call your doctor right away.
- If your sex partner is pregnant or may get pregnant. You must use a condom while taking this drug and for some time after the last dose. Use a condom even if you have had a vasectomy. Ask your doctor how long to use a condom. If your partner is pregnant or gets pregnant, call the doctor right away.
- If you are breast-feeding. Do not breast-feed while you take this drug and for 2 weeks after your last dose.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or

OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- High blood pressure has happened with this drug. Have your blood pressure checked as you have been told by your doctor.
- You may have more of a chance of getting an infection. Some infections have been severe or deadly. Wash hands often. Stay away from people with infections, colds, or flu.
- Call your doctor right away if you have any signs of infection like fever, chills, flu-like signs, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or a wound that will not heal.
- You may bleed more easily. Be careful and avoid injury. Use a soft toothbrush and an electric razor. Rarely, some bleeding problems have been deadly.
- A very bad and sometimes deadly brain problem called posterior reversible encephalopathy syndrome (PRES) has happened with this drug. Call your doctor right away if you have signs like feeling confused, lowered alertness, change in eyesight, loss of eyesight, seizures, or very bad headache.
- Very bad and sometimes deadly holes in the GI (gastrointestinal) tract or fistulas have happened with this drug. Talk with the doctor.
- Blood clots have happened with this drug. Sometimes, blood clots like heart attack and stroke have been deadly. Talk with the doctor.
- Very bad and sometimes deadly blood problems like thrombotic thrombocytopenic purpura/hemolytic uremic syndrome (TTP/HUS) have

happened with this drug in some people. Call your doctor right away if you feel very tired or weak or have any bruising or bleeding; dark urine or yellow skin or eyes; pale skin; change in the amount of urine passed; change in eyesight; change in strength on 1 side is greater than the other, trouble speaking or thinking, or change in balance; or fever.

- Severe and sometimes deadly lung problems have happened with this drug. Call your doctor right away if you have lung or breathing problems that are new or worse like cough or shortness of breath.
- Patients with cancer who take this drug may be at a greater risk of getting a severe health problem called tumor lysis syndrome (TLS). This may lead to death. Call your doctor right away if you have a fast or abnormal heartbeat; any passing out; trouble passing urine; muscle weakness or cramps; upset stomach, throwing up, diarrhea, or not able to eat; or feel sluggish.
- Rarely, severe problems with the main blood vessel from the heart (aorta) have happened with this drug. This includes aneurysm (ballooning or bulging of the aorta), tears, or bursting. This may lead to death. Call your doctor right away if you have sudden pain in the stomach, chest, or back that is severe or does not go away.
- This drug may affect fertility. Fertility problems may lead to not being able to get pregnant or father a child. If you plan to get pregnant or father a child, talk with your doctor before taking this drug.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of low thyroid levels like constipation; not able to handle cold; memory problems; mood changes; or a burning, numbness, or tingling feeling that is not normal.
- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing up.
- Chest pain or pressure.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Swelling, warmth, numbness, change of color, or pain in a leg or arm.
- Not able to pass urine or change in how much urine is passed.
- Redness or irritation of the palms of hands or soles of feet.
- Swelling of belly.
- Black, tarry, or bloody stools.
- Throwing up blood or throw up that looks like coffee grounds.
- Any unexplained bruising or bleeding.
- Heart problems like heart failure and a certain abnormal heartbeat (prolonged QT interval) have happened with this drug. Sometimes, these heart problems have been deadly. Call your doctor right away if you have a fast or abnormal heartbeat; very bad dizziness or passing out; or shortness of breath, a big weight gain, or swelling in the arms or legs.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling dizzy, tired, or weak.

- Headache.
- Upset stomach or throwing up.
- Stomach pain or diarrhea.
- Decreased appetite.
- Change in taste.
- Mouth irritation or mouth sores.
- Change in color of skin or hair.
- Hair loss.
- Muscle pain.
- Bone pain.
- Weight loss.
- Trouble sleeping.
- Change in nails.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take this drug on an empty stomach. Take at least 1 hour before or at least 2 hours after a meal.
- Swallow whole. Do not chew, break, or crush.
- Do not take antacids at the same time as this drug. Talk with your doctor.

- Avoid grapefruit and grapefruit juice.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- You may need to have an ECG checked before starting this drug and while taking it. Talk with your doctor.
- Have your urine checked as you have been told by your doctor.
- If you have upset stomach, throwing up, diarrhea, or decreased appetite, talk with your doctor. There may be ways to lower these side effects.
- This drug may affect how wounds heal. If you need to have surgery, you may need to stop this drug before surgery. Start taking it again after surgery as you have been told by your doctor. Call your doctor right away if you have a wound that does not heal or any other wound problems.
- If you are 65 or older, use this drug with care. You could have more side effects.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is less than 12 hours until the next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be

drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed

by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-02-02

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Pazopanib - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center