



---

## PATIENT & CAREGIVER EDUCATION

# How to Get Ready for Your Virtual Colonoscopy

This information will help you get ready for your virtual colonoscopy at MSK.

## About Your Virtual Colonoscopy

A virtual colonoscopy (VC) is a test to look for signs of colorectal cancer. Your doctor will look for polyps in the lining of your colon (large intestine) and rectum (the last part of your colon). Polyps are growths that usually are not cancerous but can become cancer. Your doctor will also check parts of your abdomen (belly) and pelvis for any issues.

A VC is also sometimes called a computed tomography (CT) colonography. It uses a CT scan to take 3D pictures of the inside of your colon and rectum. This helps your doctor see inside your colon without using a scope. A scope is a flexible tube placed in the rectum during a regular colonoscopy.

## Screening for colorectal cancer and polyps

Screening (checking) for colorectal cancer and polyps should start at age 45. Your doctor may recommend you screen earlier than age 45 if you have a family history of colorectal cancer. That means a close family member related to you by blood (parents, siblings, or children) has had colorectal cancer.

If your doctor finds a large polyp or growth, you may need to have a regular colonoscopy to remove it. A large polyp is 3/8 inch (1 centimeter) or bigger. Large polyps should be removed because there is a high chance they can turn into cancer.

## Before Your Procedure

### Pick up your prep kit

Before your VC, you will need to clean out (empty) your colon. To do this, you will need to pick up the following supplies **from an MSK pharmacy**:

- 2 (5 milligram) tablets of bisacodyl (Dulcolax<sup>®</sup>). You will get a box of 10 tablets, but you will only need to take 2.
- 1 (50 milliliter) bottle of iohexol 350 (Omnipaque<sup>®</sup>). This is a contrast dye that will make your colon easier to see during your VC.
  - Do not put the iohexol in the refrigerator.
  - Do not store the bottle in direct sunlight.

- 1 (225 milliliter) bottle of liquid barium sulfate suspension. This is another contrast dye.
  - You can keep the barium sulfate in the refrigerator or at room temperature.
- 1 (8.3 ounces or 238 gram) bottle of polyethylene glycol (MiraLAX<sup>®</sup>).

You will also need:

- 64 ounces of any clear liquid that isn't red, purple, or orange. You will need to mix this with MiraLAX. Keep it at room temperature.
  - A sports drink like Gatorade<sup>®</sup> or Powerade<sup>®</sup> is a good choice. Sports drinks will help replace electrolytes that you will lose during the bowel preparation.
  - If you have diabetes, be sure to get sugar-free clear liquids.
- A pitcher or bottle to mix the MiraLAX and 64 ounces of clear liquid.

## **Ask about your medications**

You may need to change the time you take your usual medications. This is so you're not taking your bowel preparation medication at the same time as your other medication. Plan to take your usual medications 1 hour before or 1 hour after taking your bowel prep medication.

## **If you're having trouble planning your medication**

**time, call your healthcare provider.**

## **Anticoagulants (blood thinners)**

Blood thinners are medications that affect the way your blood clots.

Examples of common blood thinners are listed below.

There are others, so be sure your care team knows all the medicine you take. **Do not stop taking your blood thinner without talking with a member of your care team.**

- Apixaban (Eliquis<sup>®</sup>)
- Aspirin
- Celecoxib (Celebrex<sup>®</sup>)
- Cilostazol (Pletal<sup>®</sup>)
- Clopidogrel (Plavix<sup>®</sup>)
- Dabigatran (Pradaxa<sup>®</sup>)
- Dalteparin (Fragmin<sup>®</sup>)
- Dipyridamole (Persantine<sup>®</sup>)
- Edoxaban (Savaysa<sup>®</sup>)
- Enoxaparin (Lovenox<sup>®</sup>)
- Fondaparinux (Arixtra<sup>®</sup>)
- Meloxicam (Mobic<sup>®</sup>)
- Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen (Advil<sup>®</sup>, Motrin<sup>®</sup>) and naproxen (Aleve<sup>®</sup>)
- Pentoxifylline (Trental<sup>®</sup>)
- Prasugrel (Effient<sup>®</sup>)
- Rivaroxaban (Xarelto<sup>®</sup>)
- Sulfasalazine (Azulfidine<sup>®</sup>, Sulfazine<sup>®</sup>)
- Ticagrelor (Brilinta<sup>®</sup>)

- Heparin (shot under your skin)
- Tinzaparin (Innohep<sup>®</sup>)
- Warfarin (Jantoven<sup>®</sup>, Coumadin<sup>®</sup>)

## **Contrast dye**

Contrast is a special dye that makes it easier for your doctor to see differences in the organs inside your body. You will get oral contrast to drink before your scan. If you've had a reaction (such as hives) to contrast dye in the past, tell your care team. This includes barium and iodinated contrast.

If you're having a VC without oral contrast, read *How to Get Ready for Your Virtual Colonoscopy Without Oral Contrast*

([https://sandbox18.mskcc.org/pe/vc\\_without\\_oral\\_contrast](https://sandbox18.mskcc.org/pe/vc_without_oral_contrast)).

## **Diabetes medication**

- If you take insulin or other medications for diabetes, you may need to change the dose. Ask the healthcare provider who prescribes your diabetes medication what to do the day before and morning of your procedure. Tell them you will be following a clear liquid diet the day before your procedure.
- If you monitor your blood sugar (glucose) level, ask your healthcare provider if you should stay on the same monitoring schedule. Call them if your blood sugar falls

below 70.

## **Take devices off your skin**

You may wear certain devices on your skin. Before your scan or procedure, device makers recommend you take off your:

- Continuous glucose monitor (CGM)
- Insulin pump

Talk with your healthcare provider about scheduling your appointment closer to the date you need to change your device. Make sure you have an extra device with you to put on after your scan or procedure.

You may not be sure how to manage your glucose while your device is off. If so, before your appointment, talk with the healthcare provider who manages your diabetes care.

## **3 Days Before Your Procedure**

### **Avoid certain foods**

Do not eat foods that are hard to digest. They may make it hard for your healthcare provider to see your organs in the pictures taken during your VC. You may need to have the procedure again if this happens.

Examples of foods that are hard to digest are:

- Raw fruits and vegetables. You can have vegetables that

are cooked or canned.

- Whole kernel corn, including canned corn.
- Popcorn.
- Potato skins.
- Whole grains (such as oatmeal, brown rice, quinoa, or wheat bread).
- Seeds (such as poppy or sesame).
- Nuts.

## **The Day Before Your Procedure**

### **Prepare Your MiraLAX Bowel Preparation**

On the morning of the day before your procedure, mix all 238 grams of the MiraLAX powder with 64 ounces of a room temperature clear liquid. Mix it until the MiraLAX powder dissolves. Once the MiraLAX is dissolved, you can put the mixture in the refrigerator. Many people find it tastes better when it's chilled.

Do not mix the MiraLAX earlier than the morning of the day before your procedure.

### **Start your bowel preparation**

During your bowel preparation, you will take laxatives and contrast medication. Follow these instructions carefully. It's very important that your colon is empty for your VC. If your colon isn't empty, it will be hard to see polyps or

other problems inside your colon. If this happens, you may need to do the procedure again.

If you have any questions, call your healthcare provider's office.

## **Follow a clear liquid diet**

You will need to follow a clear liquid diet the day before your procedure. A clear liquid diet includes only liquids you can see through. You can find examples in the "Clear Liquid Diet" table.

While you're following a clear liquid diet:

- Do not eat any solid foods.
- Try to drink at least 1 (8-ounce) cup of clear liquid every hour you're awake.
- Drink different types of clear liquids. Do not just drink water, coffee, and tea.
- Do not drink anything red, purple, or orange.

You can drink as many clear liquids as you like between each step of your bowel preparation. Stop drinking clear liquids at midnight.

<b>Clear Liquid Diet</b>		
	<b>OK to Drink</b>	<b>Do Not Drink</b>
<b>Soups</b>	<ul style="list-style-type: none"><li>• Clear broth, bouillon, or</li></ul>	<ul style="list-style-type: none"><li>• Any products with pieces</li></ul>



	<p>consommé.</p> <ul style="list-style-type: none"> <li>• Clear packaged vegetable, chicken, or beef broth-mix.</li> </ul>	<p>of dried food or seasoning.</p>
<b>Sweets</b>	<ul style="list-style-type: none"> <li>• Gelatin, such as Jell-O®.</li> <li>• Flavored ices.</li> <li>• Sweeteners, such as sugar or honey.</li> </ul>	<ul style="list-style-type: none"> <li>• Anything red, orange, or purple.</li> </ul>
<b>Drinks</b>	<ul style="list-style-type: none"> <li>• Water.</li> <li>• Clear fruit juices, such as white cranberry, white grape, apple.</li> <li>• Soda, such as ginger ale, 7-Up®, Sprite®, and seltzer.</li> <li>• Sports drinks, such as Gatorade®, and Powerade®.</li> <li>• Black coffee.</li> <li>• Tea.</li> <li>• Clear liquid protein drinks.</li> </ul>	<ul style="list-style-type: none"> <li>• Juice with pulp.</li> <li>• Nectars.</li> <li>• Smoothies or shakes.</li> <li>• Milk or cream.</li> <li>• Alcoholic beverages.</li> <li>• Anything red, orange, or purple.</li> </ul>

## **Follow your bowel prep medication schedule**

Start taking your bowel preparation medication the day before your VC. Follow the medication schedule in this section.

You may need to change the time you take your usual medications. If you usually take your medications at the

time you're supposed to take one of your bowel preparation medications, plan to take your usual medications 1 hour before or 1 hour after. **If you're having trouble planning your medication time, call your healthcare provider.**

If you have any questions, call 212-639-7280. If the office is closed, call 212-639-2000 and ask for the Radiology Body fellow on call.

### **Bowel prep schedule**

Use the table at the end of this resource to write down the time you take each medication. Remember that you can drink clear liquids between each of these steps. Stop drinking clear liquids at midnight.

#### **Step 1: Bisacodyl tablets**

**Anytime in the morning before 9 a.m., take 2 (5 milligram) bisacodyl tablets.**

- Take them with 1 (8-ounce) glass of clear liquids.
- Do not chew or crush them.
- Do not take them within 1 hour of taking an antacid (medication to relieve heartburn or stomach pain). Examples of antacids include calcium carbonates (such as Tums®) and histamine-2 blockers (such as Zantac®). Ask your healthcare provider for more information.

The bisacodyl tablets will make you have a bowel

movement (poop) about 6 to 8 hours after you take them. This will help the laxative you take in Step 2 and Step 3 work better. You can take these tablets and still do normal activities. They will rarely cause diarrhea (loose or watery bowel movements).

## **Step 2: First half of the MiraLAX mixture**

**At 1 p.m., drink half of the MiraLAX mixture.**

- You may want to drink it with a straw to lessen the taste.
- This is a laxative, so you should start having bowel movements more often. Your stool (poop) will become more and more loose and clear as you continue with your bowel prep. You will want to be near a bathroom. The time it takes for the laxative to start working varies for each person.
- If you do not start having bowel movements within 2 hours after completing Step 2, call your healthcare provider.
- Apply petroleum jelly (Vaseline®) or A & D® ointment to the skin around your rectum after every bowel movement. This helps prevent irritation.

Drink at least 4 to 6 cups (32 to 48 ounces) of clear liquids over the next 2 hours.

### **Step 3: Other half of the MiraLAX mixture**

**At 4 p.m., drink the other half of the MiraLAX mixture. Drink it the same way you drank the first half.**

### **Step 4: Liquid barium sulfate suspension**

**At 5 p.m., drink the bottle of liquid barium sulfate suspension.**

### **Step 5: Iohexol**

**At 7 p.m., drink the bottle of Iohexol.**

- You can mix the Iohexol with 8 ounces of clear juice, water, or soda. You can also drink the Iohexol and then follow it with 8 ounces of clear juice, water, or soda.
- The plastic bottle has a pull-tab on the top, but you do not need to pull it. Instead, twist the entire top to remove it. Take off the black rubber stopper before drinking the contrast.
- The Iohexol bottle says it's for injection, but you can also drink it. We want you to drink it so that it will be in your colon by the time of your VC.

Now your stool (poop) should be more liquidy and yellowish-clear. This is a sign that your bowels are getting cleared. If your bowels aren't getting cleared, call 212-639-7280 and ask to talk with a nurse or the fellow on call.



**Do not eat or drink anything after midnight except for your medications with a small sip of water.**

## **The Day of Your Procedure**

You may take your usual daily medications as prescribed with small sips of water.

### **What to expect**

Many staff members will ask you to say and spell your name and birth date. This is for your safety. People with the same or a similar name may be having surgery on the same day.

When it's time to change for your procedure, you will get a hospital gown to wear.

You will be taken into the CT exam room where you will lie on the CT exam table. A small tube will be gently placed into your rectum and secured to your leg or buttocks. It will feel like you're having a rectal exam. The tube will inflate your colon with air or gas so that your technologist can see inside it. Your technologist will ask you to change positions a few times. This is to help move the air or gas throughout your colon. You will lie on your

side, back, and abdomen (belly).

The VC is not painful, but you may feel some bloating, discomfort, or cramping. You may also feel like you're going to have a bowel movement. These feelings should go away after your procedure.

Your technologist will ask you to hold your breath for about 10 seconds while the CT scanner takes pictures. They will take pictures of your abdomen and pelvis in at least 2 different positions. This is usually while you're on your back and stomach.

## **After Your Procedure**

- You may go back to your normal diet after your procedure.
- You may feel some cramping in your stomach. This is because of the gas used during your procedure. Walking can help relieve gas and reduce your cramping.
- It usually takes 2 to 3 business days (Monday through Friday) to get your results. The doctor who sent you for your VC will call you when your results are ready.

## **When to Call Your Healthcare Provider**

Call your healthcare if you have:

- A fever of 101° F (38.4° C) or higher.
- Stomach pain, other than mild cramping.
- Bleeding from your rectum, A small amount of blood on toilet paper is normal 24 hours after your VC.
- Weakness or faintness.
- Nausea (feeling like you're going to throw up).
- Vomiting (throwing up).

## Bowel Prep Schedule

Use this table to write down the time you take each medication.

Step	Scheduled Time	Actual Time Taken	Medication(s) to Take
1	Before 9 a.m.		<ul style="list-style-type: none"> <li>• Take 2 (5 milligram) bisacodyl tablets.</li> </ul>
2	At 1 p.m.		<ul style="list-style-type: none"> <li>• Drink half of the MiraLAX mixture (32 ounces).</li> </ul>
3	At 4 p.m.		<ul style="list-style-type: none"> <li>• Drink the other half of the MiraLAX mixture (32 ounces).</li> </ul>

<b>4</b>	At 5 p.m.		<ul style="list-style-type: none"> <li>• Drink 1 (225 milliliter) bottle of liquid barium sulfate suspension.</li> </ul>
<b>5</b>	At 7 p.m.		<ul style="list-style-type: none"> <li>• Drink 1 (50 milliliter) bottle of iohexol.</li> <li>• You can mix this with an 8 ounce clear liquid or drink the iohexol and then drink an 8 ounce clear liquid.</li> </ul>

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

How to Get Ready for Your Virtual Colonoscopy - Last updated on February 13, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center